


Nitrogen Production and Excretion in Fish

David J. Randall
Don D. MacKinlay



International Congress on the Biology of Fish
Towson University, Baltimore MD July 26-30, 1998

***Nitrogen Production and
Excretion in Fish***

SYMPOSIUM PROCEEDINGS

Dave Randall

Don MacKinlay

*International Congress on the Biology of Fish
Towson University, Baltimore MD July 27-30, 1998.*

Copyright © 1998
Physiology Section,
American Fisheries Society
All rights reserved

International Standard Book Number (ISBN)1-894337-03-4

Notice

This publication is made up of a combination of extended abstracts and full papers, submitted by the authors without peer review. The papers in this volume should not be cited as primary literature. The Physiology Section of the American Fisheries Society offers this compilation of papers in the interests of information exchange only, and makes no claim as to the validity of the conclusions or recommendations presented in the papers.

For copies of these Symposium Proceedings, or the other 20 Proceedings in the Congress series, contact:

Don MacKinlay, SEP DFO, 555 West Hastings St.,
Vancouver BC V6B 5G3 Canada
Phone: 604-666-3520 Fax 604-666-6894
E-mail: mackinlayd@pac.dfo-mpo.gc.ca

Website: www.fishbiologycongress.org

PREFACE

Teleosts are the major vertebrate group with between 20,000 and 30,000 species representing about 50 to 60% of all vertebrate species. Vertebrate movement onto land is a relatively recent event and much of the basic design of vertebrates was probably established in the aquatic environment. The broad generalization that aquatic vertebrates are ammonotelic, whereas terrestrial vertebrates are ureotelic, has obscured the fact that most mechanisms of nitrogen excretion are found in all vertebrate classes and that these mechanisms probably evolved in vertebrate aquatic ancestors. Teleosts and present day land vertebrates have diverged considerably from the ancestral aquatic vertebrate stock, but comparative studies of all vertebrates may elucidate basic patterns in nitrogen excretion in vertebrates.

Ammonia is continually produced by all vertebrates but cannot be accumulated in the body because it is toxic. Ammonia must, therefore, be excreted or converted to some other less toxic compound. Fish have evolved a wide variety of mechanisms to avoid ammonia toxicity, the symposium on NITROGEN PRODUCTION AND EXCRETION IN FISH describes and discusses many of these mechanisms. There is also some discussion of the sub-lethal toxic action of ammonia. In addition there is a review of ammonia transport in the mammalian nephron, with discussion of some recent developments in that field.

Dave Randall
University of B.C

Don MacKinlay
Habitat & Enhancement Branch

CONGRESS ACKNOWLEDGEMENTS

This Symposium is part of the International Congress on the Biology of Fishes, whose main sponsors were Fisheries and Oceans Canada (DFO), and Towson University. The main organizers of the Congress, on behalf of the Physiology and Fish Culture Sections of the American Fisheries Society, were Don MacKinlay of DFO (overall chair, program and proceedings), Karin Howard (registration and accommodations) and Jay Nelson of Towson University (local arrangements). I would like to extend a sincere 'thank you' to the many contributors who took the time to prepare a written submission for these proceedings. Your efforts are very much appreciated.

Don MacKinlay
Congress Chair

TABLE OF CONTENTS

Urea synthesis/excretion and ammonia toxicity in the gulf toadfish (<i>Opsanus beta</i>) <i>Walsh, P.J., Y. Wang & C.M. Wood</i>	5
Urea metabolism and excretion in the Lake	5
A.	5
nephron and its glucocorticoids.
Ammonia excretion in the <i>Ip</i> , <i>L.N.Hong, T.W.K.</i> , <i>Peng,</i>	15
<i>Wilson, R.W. & D.L.</i> acidophilic Amazonian	19
exposed to copper in soft acidic water.	27
lamprey (<i>marinus</i> <i>Wilkie,</i> <i>J. Holmes & J.H.</i>	31
sublethal un-ionized tilapia <i>Oreochromis niloticus</i> <i>Abdalla, A.A.F. & C.D.</i>	35
Inari charr <i>Salvelinus alpinus</i> <i>Lyytikainen, T. & M.</i>	49

Expression of glutamine synthetase and ornithine carbamoyl transferase in the course of <i>Piaractus mesopotamicus</i> Holberg 1887 (Teleostei, Characidae) development. <i>Monzani, P.S., S.M. Ramos, R.O. Ramos & G. Moraes</i>	55
Nitrogenous excretion pattern of the fresh water fish <i>Hoplerytinus unitaeniatus</i> (jeju): a facultative air-breathing neotropical teleost. <i>Polez, V.L.P., G. Moraes & R. Huet de Salvo Souza</i>	63